Needle Exchange Technology (NEXT): A novel internet-based maildelivered syringe services program

BENJAMIN T HAYES, MD, MSW, MPH JAMIE FAVARO, EXECUTIVE DIRECTOR, NEXT DISTRO

OCTOBER 7, 2020







Albert Einstein College of Medicine





Conflicts of interest

- Benjamin Hayes has no conflicts of interest to declare.
- Jamie Favaro is Executive Director of NEXT; no other conflicts to declare.



Background

- Most people in the US have poor access to syringe services programs (SSPs)
 - > Rural areas have far fewer SSPs than urban areas¹
 - > Women and young people are often underrepresented in SSP programs
 - Stigma, fear of legal consequences and geography act as major barriers to inperson SSPs ^{2, 3, 4, 5, 6, 7}



93% of the 220 counties most vulnerable to HIV or HCV outbreaks are without SSP^{8,9}



15 vulnerable counties have SSPs

NEXT DISTRO & NEXT NALOXONE

- NEXT DISTRO (began distributing 2018)
 - sterile injection equipment and other harm reduction supplies
- NEXT NALOXONE
 - naloxone for overdose prevention
- Goal: augment existing SSPs by
 - 1. Supporting people not otherwise accessing in-person SSPs
 - 2. Helping local services expand their reach
- Internet-based technology \rightarrow mail distribution
- Encrypted messaging service (confidentiality)
- First formal program in US



How does it work:



Getting Sterile Injection Equipment:



Getting Harm Reduction Supplies:



NY NEXT

CA NEXT



OTHER STATES





Getting Naloxone:

REVERSING AN OPIOID OVERDOSE IS

IN YOUR HANDS

Get Naloxone

Report Naloxone Use

ALOXONE HIS ALOXONE HIS ALOXONE HIS ACTION, USP 0.4 mg IV, I.M., or S.C. one mot from light. Rushing the second second second second motor from the second second second second motor from the second motor from the second se

4 mg

icalcunne HCL

NARCAN® NASAL SPRAY

HE NOSE ON

ALABAMA	INDIANA	NEBRASKA	PUERTO RICO
ALASKA	IOWA	NEVADA	RHODE
ARIZONA	KANSAS	NEW HAMPSHIRE	ISLAND
ARKANSAS	KENTUCKY	NEW JERSEY	SOUTH
CALIFORNIA	LOUISIANA	NEW MEXICO	CAROLINA
COLORADO	MAINE	NEW YORK	SOUTH
CONNECTICUT	MARYLAND	NORTH CAROLINA	DAKOTA
	MASSACHUSETTS	NORTH DAKOTA	TENNESSEE
DELAWARE	MICHIGAN	ОНЮ	TEXAS
FLORIDA	MINNESOTA	OKLAHOMA	UTAH
GEORGIA	MISSISSIPPI	OREGON	VERMONT
HAWAI'I	MISSOURI	PENNSYLVANIA	VIRGINIA
IDAHO	MONTANA	PHILADELPHIA	WASHINGTON
ILLINOIS			WEST
			VIRGINIA
			WISCONSIN

WYOMING

		NERDACKA	
ALADAMA	INDIANA	NEDRASKA	PUERIO RICO
ALASKA	IOWA	NEVADA	RHODE
ARIZONA	KANSAS	NEW HAMPSHIRE	ISLAND
ARKANSAS	KENTUCKY	NEW JERSEY	SOUTH
CALIFORNIA	LOUISIANA	NEW MEXICO	CAROLINA
COLORADO	MAINE	NEW YORK	SOUTH
CONNECTICUT	MARYLAND	NORTH CAROLINA	DAKOTA
	MASSACHUSETTS	NORTH DAKOTA	TENNESSEE
DELAWARE	MICHIGAN	OHIO	TEXAS
FLORIDA	MINNESOTA	OKLAHOMA	UTAH
GEORGIA	MISSISSIPPI	OREGON	VERMONT
HAWAI'I	MISSOURI	PENNSYLVANIA	VIRGINIA
IDAHO	MONTANA	PHILADELPHIA	WASHINGTON
ILLINOIS			WEST
			VIRGINIA
			WISCONSIN
NG >			WYOMING



RESOURCES IN

North Carolina

In 2018, there were 1,783 opioid overdose deaths in the state of North Carolina.

If you are a person who uses drugs or if you are around people who use drugs, please consider learning how to respond to an opioid overdose and carrying naloxone (intranasal brand name: Narcan). Naloxone is a medication designed to reverse an opioid overdose. Opioids slow down the activity of the central nervous system. This can cause slowed or stopped breathing which may lead to death. Naloxone displaces the opioids from the opioid receptors in the brain which allows the person who is experiencing an opioid overdose to begin breathing normally. We have a helpful FAQ and are constantly updating it with questions from our site visitors. If you don't find what you're looking for, let us know.













Naloxone Finder

Good Samaritan Policy

Crisis Support

Syringe Access

Pharmacy

Organizing

GET NALOXONE MAILED TO YOU

If you need naloxone and **cannot access it locally**, you may qualify to have intramuscular naloxone mailed to you.







NEXT DISTRO: sterile injection supplies program

• Who does the program serve?

Methods:

- Participants: newly requesting supplies 2/18 8/19
- Data collection: self-report enrollment form from website
- Measures: sociodemographic info, self-reported overdose, HCV infection, previous source of supplies



Results

Participant characteristics	N = 173 (%)
Mean age	35 yrs (20 – 64 yrs); 50% <33 yrs
White Latinx Black or African American Other	87% 6% 1.7% 5.2%
Female	53%
Unstable housing or homeless	41%
Uninsured	59%

Results

Drugs used past 30 days: Heroin Methamphetamine Opioid analgesics	69% 51% 34%
Overdoses, past 12 months	25%
Witnessed overdose in past 12 months	66%
HCV infection, self-report	38% 50% of women
Distributing supplies to other people	63% (19 missing)

Previous Sources of Injection Supplies





- High frequency of overdose and HCV infection
- Few participants previously used pharmacies or SSPs for supplies
- NEXT reaches large proportion of women and young adults, which are often underrepresented in traditional SSPs



Next steps

• Research:

- Evaluate NEXT's data on services to underserved populations
- Evaluate NEXT Naloxone services
- NEXT service delivery:
 - Building their own encrypted web application
 - Other SSPs will be able to use this app to expand their services
 - Evolve the website into an open-access, multilanguage hub for drug user resources
 - Expand to offer more telehealth services



Acknowledgements

- Thank you to NEXT for their collaboration.
- Thank you to Dr. Aaron D. Fox for his mentorship and to Dr Czarina Behrends for her insights and suggestions.

References

1. Jarlais, D. C. D., Nugent, A., Solberg, A., Feelemyer, J., Mermin, J., & Holtzman, D. (2015). Syringe service programs for persons who inject drugs in urban, suburban, and rural areas—United States, 2013. *Morbidity and mortality weekly report*, *64*(48), 1337-1341.

3. Beletsky L, Heller D, Jenness SM, Neaigus A, Gelpi-Acosta C, Hagan H. Syringe access, syringe sharing, and police encounters among people who inject drugs in New York City: a community-level perspective. *Int J Drug Policy*. Jan 2014;25(1):105-11. doi:10.1016/j.drugpo.2013.06.005

2. Rivera AV, DeCuir J, Crawford ND, Amesty S, Lewis CF. Internalized stigma and sterile syringe use among people who inject drugs in New York City, 2010-2012. *Drug Alcohol Depend*. Nov 1 2014;144:259-64. doi:10.1016/j.drugalcdep.2014.09.778

4. Cloud, D. H., Ibragimov, U., Prood, N., Young, A. M., & Cooper, H. L. (2019). Rural risk environments for hepatitis c among young adults in appalachian kentucky. *International Journal of Drug Policy*, 72, 47-54.

5. Delany-Moretlwe, S., Cowan, F. M., Busza, J., Bolton-Moore, C., Kelley, K., & Fairlie, L. (2015). Providing comprehensive health services for young key populations: needs, barriers and gaps. *Journal of the International AIDS Society*, *18*, 19833.

6. El-Bassel, N., & Strathdee, S. A. (2015). Women who use or inject drugs: an action agenda for women-specific, multilevel and combination HIV prevention and research. *Journal of acquired immune deficiency syndromes (1999), 69*(Suppl 2), S182.

7. Canary, L., Hariri, S., Campbell, C., Young, R., Whitcomb, J., Kaufman, H., & Vellozzi, C. (2017). Geographic disparities in access to syringe services programs among young persons with hepatitis C virus infection in the United States. *Clinical Infectious Diseases*, *65*(3), 514-517.

8. amFAR. (2017, November 3). Syringe Services Programs and the Opioid Epidemic. Retrieved August 26, 2020, from https://www.amfar.org/ssp-opioid-epidemic/

9. Van Handel MM, Rose CE, Hallisey EJ, et. al. County-Level Vulnerability Assessment for Rapid Dissemination of HIV or HCV Infections Among Persons Who Inject Drugs, United States. J Acquir Immune Defic Syndr. 2016 Nov 1;73(3): 323-331 <u>https://www.cdc.gov/pwid/vulnerable-counties-data.html</u>